

CORRECTIONS

In the Original Article entitled “Association of intrarenal blood flow with renal function and target organ damage in hypertensive patients with fibromuscular dysplasia: the ARCADIA-POL study” published in the April 2019 issue (Januszewicz M, Januszewicz A, Michałowska I, et al. *Pol Arch Intern Med.* 2019; 129 (4): 234–241). doi:10.20452/pamw.4479) **TABLE 4** was omitted. The table was added on page 241 of the article, as below.

TABLE 4 Renal arteries with no fibromuscular dysplasia lesions, fibromuscular dysplasia and nonsignificant renal artery stenosis, fibromuscular dysplasia and significant renal artery stenosis

Parameter	Renal arteries without FMD	Renal arteries with FMD	Renal arteries with FMD	P value			
				ANOVA	1 vs 2	1 vs 3	2 vs 3
Length of kidney, mm	112.4 (10.1)	108.4 (10.7)	107.4 (8.2)	0.05	0.09	0.14	1.00
V _{max} , cm/s	116.0 (43.8)	151.7 (57.7)	314.8 (105.6)	<0.001	<0.001	<0.001	<0.001
RAR	1.18 (0.47)	1.89 (0.75)	4.19 (2.09)	<0.001	0.005	<0.001	<0.001
RRI	0.60 (0.06)	0.59 (0.07)	0.47 (0.09)	<0.001	0.50	<0.001	<0.001
AT	20.2 (7.6)	24.2 (12.2)	61.4 (32.0)	<0.001	0.16	<0.001	<0.001
PI	1.08 (0.18)	1.09 (0.19)	1.12 (0.22)	0.64	1.00	1.00	1.00

Data are presented as mean (SD).

Abbreviations: AT, acceleration time; PI, pulsatility index; RAR, renal aortic ratio; V_{max}, maximal blood flow velocity; others, see **TABLE 1**

The article is correct at www.mp.pl/paim.