

Pitfalls in the management of bilateral adrenal enlargement

Piotr Wiśniewski¹, Łukasz Obołończyk¹, Marta Gawrońska²,
Dominika Okrój², Piotr Czapiewski³, Krzysztof Sworczak¹

¹ Department of Endocrinology and Internal Medicine, Medical University of Gdańsk, Gdańsk, Poland

² Endocrinology Student's Scientific Group, Medical University of Gdańsk, Gdańsk, Poland

³ Department of Pathomorphology, Medical University of Gdańsk, Gdańsk, Poland

Unsuspected adrenal masses, or incidentalomas, are increasingly found with the widespread use of thoracic and abdominal imaging; most cases do not pose a significant risk to a patient's health.¹ We describe a case of a patient with enlarged adrenals who was initially diagnosed with nonclassic congenital adrenal hyperplasia (NCAH), but 1 year later died from primary adrenal lymphoma dissemination.

In January 2013, an 83-year-old woman visited a general practitioner due to nonspecific chronic abdominal pain. Abdominal ultrasound suggested an adrenal mass and computed tomography (CT) scanning revealed bilateral adrenal enlargement. The glands measured up to 69 mm and had a baseline density of 38 HU and contrast wash-out rate below 60% (FIGURE 1A). No other abdominal lesions were detected.

In April 2013, the patient was admitted to our hospital for diagnostic workup. Physical

examination was unremarkable except for mild abdominal tenderness. Routine laboratory test results were normal. The circadian cortisol profile, urinary cortisol excretion, low-dose dexamethasone suppression test results, as well as serum testosterone and androstenedione levels were normal. The baseline and stimulated 17-hydroxyprogesterone levels were increased to 6.94 ng/ml and 20 ng/ml, respectively.

The patient was diagnosed with NCAH and received no treatment. Because of the indeterminate CT features, adrenal biopsy was considered but the patient refused.

In September 2013, she presented with classic symptoms of adrenal crisis, and steroid replacement was initiated. Adrenal CT was ordered; however, due to the patient's personal reasons, the scan was performed as late as in April 2014. It revealed enormous adrenals measuring up to 156 mm (FIGURE 1B) with periadrenal fat tissue infiltration.




FIGURE 1 Computed tomography (CT) imaging: **A** – an abdominal CT scan performed in January 2013, showing bilateral enlargement of the adrenal glands (arrows). The phenotype was not specific for lipid-rich adenoma. **B** – a follow-up CT scan performed in April 2014 and showing a marked progression of adrenal enlargement (arrows)

Correspondence to:
Piotr Wiśniewski, MD, PhD,
Katedra i Klinika Endokrynologii
i Chorób Wewnętrznych,
Uniwersyteckie Centrum Kliniczne,
ul. Dębinki 7, 80-952 Gdańsk,
phone: +48 58 349 28 40, email:
piotr.wisniewski@gumed.edu.pl
Received: October 13, 2017.
Revision accepted:
October 30, 2017.
Published online:
December 21, 2017.
Conflict of interests: none declared.
Pol Arch Intern Med. 2017;
127 (12): 887-888
doi:10.20452/pamw.4175
Copyright by Medycyna Praktyczna,
Kraków 2017






FIGURE 1 Histopathological examination; **C** – dense infiltrate of neoplastic cells showing a high mitotic rate, open chromatin with multiple nucleoli, sparse cytoplasm, and numerous apoptotic bodies; **D** – strong diffuse expression of CD20; **E** – high Ki-67 proliferation index (nearly 100%)



Laboratory tests revealed elevated lactate dehydrogenase and β_2 -microglobulin in serum (1076 U/l and 8.6 mg/l, respectively). A core needle biopsy of the adrenal gland was performed, and a histopathological examination confirmed diffuse large B-cell lymphoma (DLBCL, **FIGURE 1C–1E**). The patient died before receiving the first dose of chemotherapy due to severe infectious complications. On autopsy, bilateral adrenal infiltration as well as involvement of the kidney, liver, stomach, peritoneum, and multiple periaortic lymph nodes were found.

In most cases, bilateral adrenal masses represent a benign adrenocortical disease: bilateral adenomas, macronodular hyperplasia, or distinct bilateral nodules with normal or atrophic cortex intervening.² A differential diagnosis includes metastases, adrenal hemorrhage, some infections (eg, tuberculosis), and lymphoma.³

Current guidelines recommend a determination of whether an adrenal mass can be classified as a lipid-rich adenoma based on CT or magnetic resonance imaging.^{2,4} Other types of tumors increase the likelihood of malignancy and require in-depth diagnostic workup, careful follow-up, or surgery.

The initial CT findings in our patient were not consistent with adrenal adenoma. However, given the patient's asymptomatic presentation and the hormonal test results, NCAH was diagnosed. The diagnosis was reconsidered after the patient developed adrenal crisis, and because of her poor compliance, the control CT and the resulting biopsy were delayed. Rapid tumor growth could have been probably detected by any suitable imaging modality, including ultrasound. The point-of-care ultrasound examination performed at the time

of adrenal crisis could lead to a more timely diagnosis of adrenal DLBCL.

Although most patients with adrenal tumors have good prognosis, a thorough initial evaluation and periodic follow-up are necessary to identify those at risk for poor outcome. The presented case underscores the importance of the recently published guidelines concerning adrenal tumors.⁴

REFERENCES

- 1 Young WF. The incidentally discovered adrenal mass. *N Engl J Med*. 2007; 356: 601-610.
- 2 Fassnacht M, Arit V, Bancos I, et al. Management of adrenal incidentalomas: European Society of Endocrinology Clinical Practice Guideline in collaboration with the European Network for the Study of Adrenal Tumors. *Eur J Endocrinol*. 2016; 175: G1-G34.
- 3 Kasperlik-Zaluska AA, Słowińska-Szednicka J, Rosłonowska E, et al. Bilateral, incidentally found adrenal tumours – results of observation of 1790 patients registered at a single endocrinological centre. *Endokrynol Pol*. 2010; 61: 69-73.
- 4 Bednarczuk T, Bolański M, Sworczak K, et al. Adrenal incidentaloma in adults – management recommendations by the Polish Society of Endocrinology. *Endokrynol Pol*. 2016; 67: 234-258.