

Supplementary material

Chomicka I, Kwiatkowska M, Małyszko J. Post-contrast acute kidney injury following computed tomography: a real or overestimated threat? Pol Arch Intern Med. 2020; 130: 704-707. doi:10.20452/pamw.15409

Please note that the journal is not responsible for the scientific accuracy or functionality of any supplementary material submitted by the authors. Any queries (except missing content) should be directed to the corresponding author of the article.

Table S1

Basal clinical and biochemical characteristics of patients undergoing computed tomography (CT)

Parameter:	CT with contrast (n=123)	CT without contrast (n=29)	P value
Age, years	67 (19)	63 (25)	0.87
Gender, male/female, %	51 /49	51 /49	0.83
Mode, urgent %	45	39	0.67
Mode, elective, %	55	61	0.63
AKI before CT, %	16	38	0.037
CKD, %	21	38	0.048
DM, %	25	17	0.51
Hypertension, %	54	59	0.68
Cancer, %	24	21	0.81
CHF, %	24	35	0.40
CAD, %	18	21	0.82
Liver Failure, %	7	3	0.041
Sepsis/severe infection, %	22	48	0.03
Anemia, %	54	72	0.048
Mean Hgb, mmol/L	7.45 (3.72)	6.89 (1.55)	0.29

Data given are mean (SD) or percentage

Abbreviations: AKI, acute kidney injury; CAD, coronary artery disease; CHF, chronic cardiac failure; CKD, chronic kidney disease; CT, computed tomography; DM, diabetes mellitus; eGFR, estimated glomerular filtration rate; Hgb, hemoglobin,