

Supplementary material

Zeng F, Li L, Zeng J, et al. Can we predict the severity of coronavirus disease 2019 with a routine blood test? Pol Arch Intern Med. 2020; 130: 400-406.
doi:10.20452/pamw.15331

Please note that the journal is not responsible for the scientific accuracy or functionality of any supplementary material submitted by the authors. Any queries (except missing content) should be directed to the corresponding author of the article.

Table S1. Newcastle–Ottawa Scale (NOS) score of included studies

Included studies	Year	Is the definition adequate?	Representativeness of the cases	Selection of controls	Definition of controls	Comparability of both groups	Ascertainment of diagnosis	Same ascertainment method for both groups	Nonresponse rate	Total scores
Chen L	2020	☆	☆	☆	☆	-	☆	☆	☆	7
Chen X	2020	☆	☆	☆	☆	-	☆	☆	☆	7
Cheng K	2020	☆	☆	☆	☆	☆	☆	☆	☆	8
Fang X	2020	☆	☆	☆	☆	-	☆	☆	☆	7
Gao Y	2020	☆	☆	☆	☆	☆☆	☆	☆	☆	9
Guan W	2020	☆	☆	☆	☆	☆	☆	☆	☆	8
Li K	2020	☆	☆	☆	☆	-	☆	☆	☆	7
Liu M	2020	☆	☆	☆	☆	-	☆	☆	☆	7
Peng Y	2020	☆	☆	☆	☆	☆	☆	☆	☆	8
Qin C	2020	☆	☆	☆	☆	-	☆	☆	☆	7
Qu R	2020	☆	☆	☆	☆	-	☆	☆	☆	7
Wu C	2020	☆	☆	☆	☆	-	☆	☆	☆	7
Xiang T	2020	☆	☆	☆	☆	☆	☆	☆	☆	8
Xiao K	2020	☆	☆	☆	☆	-	☆	☆	☆	7
Zhang J	2020	☆	☆	☆	☆	-	☆	☆	☆	7




Figure S1. Sensitivity analyses between non-severe and severe groups. A. White blood cell (WBC); B. Neutrophils; C. Lymphocytes; D. Neutrophil-to-lymphocyte ratio (NLR); E. Monocytes; F. Platelets.




Figure S2. Publication bias between non-severe and severe groups by Egger's test. A. White blood cell (WBC); B. Neutrophils; C. Lymphocytes; D. Neutrophil-to-lymphocyte ratio (NLR); E. Monocytes; F. Platelets.