

## Supplementary material

Majdan A, Majdan M, Dryglewska M, et al. The presence of particular criteria and noncriteria antiphospholipid antibodies in patients with uterine malignancies. *Pol Arch Intern Med.* 2020; 130: 1037-1042. doi:10.20452/pamw.15624

Please note that the journal is not responsible for the scientific accuracy or functionality of any supplementary material submitted by the authors. Any queries (except missing content) should be directed to the corresponding author of the article.

## Supplementary material

Table S1. Patients characteristics - postoperative diagnoses in group with uterine malignancies.

Postoperative diagnosis	Number of patients	Age years: mean and standard deviation (SD)	Clinical Stage FIGO (%)	Grading (%)
<i>Adenocarcinoma endometrialis</i>	68	60.4 (12.85)	I - 43 (63.2%) II - 11 (16.2%) III - 10 (14.7%) IV - 2 (2.94%) No data - 2 (2.94%)	G1 - 6 (8,8%) G2 - 51 (75%) G3 - 8 (11.8%) No data - 3 (4,4%)
<i>Leiomyosarcoma</i>	2	62 (9.14)	I - 1 (50%) No data - 1	G1 - 1 (50%) No data - 1

			(50%)	(50%)
--	--	--	-------	-------

Table S2. Patients characteristics: postoperative diagnoses in group with non - cancerous lesions of female genital tract.

<b>Postoperative diagnosis</b>	<b>Number of patients</b>	<b>Age years: mean and standard deviation (SD)</b>
<i>Endometriosis</i>	25	39.24 (10.60)
<i>Cystis simplex ovarii</i>	19	46.17 (13.65)
<i>Cystis dermoidalis</i>	10	32.30 (6.68)
<i>Leiomyoma</i>	8	52.75 (9.44)
<i>Cystadenoma mucinosum</i>	4	45.5 (22.34)
<i>Cystadenoma serosum</i>	3	58 (3.61)
<i>Fibrothecoma</i>	3	67.67 (27.79)
<i>Salpingitis</i>	2	32.5 (9.19)
<i>Endometrium involutivum</i>	2	64.5 (3.54)
<i>Polypus endometrialis</i>	2	62 (3.21)
<i>Cystadenoma multiloculare</i>	1	55
<i>Hyperplasia endometrii simplex</i>	1	64
<i>Benign mesenchymoma</i>	1	47

