

Supplementary material

Wójcik K, Masiak A, Jeleniewicz R, et al; POLVAS consortium. Association of antineutrophil cytoplasmic antibody (ANCA) specificity with demographic and clinical characteristics of patients with ANCA-associated vasculitides. *Pol Arch Intern Med.* 2022; 132: 16187. doi:10.20452/pamw.16187

Please note that the journal is not responsible for the scientific accuracy or functionality of any supplementary material submitted by the authors. Any queries (except missing content) should be directed to the corresponding author of the article.

Supplementary Table 1. Clinical characteristics and organ involvement in ANCA-negative AAV subjects.

	ANCA-negative		
	MPA	GPA	EGPA
Numbers of total AAV	2	26	47
Organ involvement			
General n (%)	1(50)	20(76.9)	46(97.9)
Skin n (%)	2(100)	8(30.8)	25(53.2)
Eye n (%)	0(0)	12(46.1)	1(2.1)
ENT n (%)	0(0)	25(96.1)	38(80.8)
Respiratory n (%)	1(50)	13(50.0)	47(100)
Heart n (%)	0(0)	6(23.1)	25(53.2)
GI n (%)	0(0)	0(0)	15(31.9)
Renal n (%)	2(100)	3(11.5)	10(21.3)
Urinary n (%)	0(0)	1(3.8)	0(0)
CNS n (%)	0(0)	2(7.7)	4(8.5)
Neurological n (%)	0(0)	3(11.5)	24(51.1)
Creatinine max (mg/l); median; [IQR]	2.36	1.1 [0.9 -2]	0.81 [0.7-0.99]
CRP at baseline (mg/l); median; [IQR]	22	9 [2-28]	30.5 [12-57]
Number of organs involved; median [IQR]	2.5	3[2-3.75]	4 [3-5]
Relapse n (%)	0(0)	15(57.7)	35(74.5)
Dialysis n (%)	0(0)	0(0)	0(0)
Dialysis transient n (%)	0(0)	0(0)	0(0)