

Supplementary material





Terlecki M, Kocowska-Trytko M, Plens K, et al. Prognostic value of acid-base balance parameters assessed on admission in peripheral venous blood of patients with myocardial infarction treated with percutaneous coronary intervention. Pol Arch Intern Med. 2022; 132: 16229.

doi:10.20452/pamw.16229

Please note that the journal is not responsible for the scientific accuracy or functionality of any supplementary material submitted by the authors. Any queries (except missing content) should be directed to the corresponding author of the article.





Figure S1 Correlation between lactate, anion gap, base excess and troponin concentration on admission and Global Registry of Acute Coronary Events (GRACE) score. Whole study group (A) and subgroup of patients without out-of-hospital cardiac arrest (B)

A.



Variable	By Variable	Spearman's correlation	Spearman's correlation P-value
GRACE score	Lactates (mmol/l)	0.361	<0.001
GRACE score	Anion Gap (mEq/l)	0.158	0.004
GRACE score	Base Excess (mEq/l)	-0.383	<0.001
GRACE score	Troponin I (ug/l)	0.097	0.079

B.



Variable	By Variable	Spearman's correlation	Spearman's correlation P-value
GRACE score	Lactates (mmol/l)	0.253	<0.001
GRACE score	Anion Gap (mEq/l)	0.076	0.180
GRACE score	Base Excess (mEq/l)	0.297	<0.001
GRACE score	Troponin I (ug/l)	0.108	0.057