

Supplementary material

Dobrowolska K, Brzdęk M, Zarębska-Michaluk D, et al. Differences between the course of SARS-CoV-2 infections in the periods of the Delta and Omicron variant dominance in Poland. Pol Arch Intern Med. 2023; 133: 16403. doi:10.20452/pamw.16403

Please note that the journal is not responsible for the scientific accuracy or functionality of any supplementary material submitted by the authors. Any queries (except missing content) should be directed to the corresponding author of the article.

Table S1. Frequency of outcomes in patients hospitalized during periods of Delta and Omicron variant dominance and treated with or without antivirals

Parameters	Delta- All, n=1440	Omicron- All, n=785	<i>P</i>	Delta- Females, n=687	Omicron- Females, n=407	<i>P</i>	Delta- Males, n=753	Omicron- Males, n=378
Frequency of events with use of antivirals/without use of antivirals % (n)								
Mortality	11.1 (52)/ 16.6 (162) p=0.01	9.9 (40)/ 16.4 (62) p=0.01	0.54/ 0.90	11.4 (25)/ 18.6 (87) p=0.02	10.2 (21)/ 14.4 (29) p=0.19	0.69/ 0.19	10.9 (27)/ 14.9 (75) p=0.13	9.5 (19)/ 18.5 (33) p=0.01
Mechanical ventilation	7.5 (35)/ 7 (68) p=0.73	2.5 (10)/ 3.7 (14) p=0.32	<0.001/ 0.02	7.8 (17)/ 6 (28) p=0.38	2.9 (6)/ 2 (4) p=0.55	0.03/ 0.03	7.3 (18)/ 7.9 (40) p=0.75	2 (4)/ 5.6 (10) p=0.06