Bednarkiewicz J, Grzejszczak P, Szmigielska-Kapłon A, et al. Acute kidney injury in a patient with T-cell acute lymphoblastic leukemia. Pol Arch Intern Med. 2023; 133: 16346. doi:10.20452/pamw.16346

Please note that the journal is not responsible for the scientific accuracy or functionality of any supplementary material submitted by the authors. Any queries (except missing content) should be directed to the corresponding author of the article.

Supplementary Table S1. The patient treatment according to PALG-7 protocol (Polish Adult Leukemia Study Group) prior to transplantation.

Medication	Dose	Time
Induction treatment		
Dexamethason orally	40 mg daily	2 days weekly
Vincristine iv	2 mg	1 dose weekly, totally 4
		doses
Daunorubicin iv	50 mg/m^2	1 dose weekly, totally 4
		doses
PEG-asparaginase	2000 IE/ m^2	1 dose, day 20
Consolidation I started at 6 week of treatment,		
Methotrexate iv	$1,5 \text{ g/m}^2$	Day 1 and 14
Etoposide iv	$100 \text{ mg} / \text{m}^2$	Day 1 and 14
Consolidation II started at 10 week of treatment		
Cytarabine iv	$2x2g/m^2$	Day 1 and 18
Cyclophosphamide iv	$1g/m^2$	Day 2-3 and 19-20
PEG-asparaginase	2000 IE/ m^2	Day 15 and 22

• Consolidation treatment is given after confirmation of complete remission of the disease. The patient received induction and both consolidation courses and then proceeded for transplantation with Cyclophosphamide/TBI (Total Body Irradiation) conditioning.