

Supplementary material

Łoń I, Leśnowolski Ł, Janiec I, et al. Severe course of type IV Takayasu disease during treatment with etanercept. *Pol Arch Intern Med.* 2024; 134: 16606.
doi:10.20452/pamw.16606

Please note that the journal is not responsible for the scientific accuracy or functionality of any supplementary material submitted by the authors. Any queries (except missing content) should be directed to the corresponding author of the article.

The patient's score in accordance with ACR/EULAR classification criteria for the diagnosis of Takayasu arteritis¹

Absolute requirements	
Age ≤ 60 years at time of diagnosis	+
Evidence of vasculitis on imaging	+
Additional clinical criteria	
Female sex	+1
Angina or ischemic cardiac pain	0
Arm or leg claudication	0
Vascular bruit	+2
Reduced pulse in upper extremity	0
Carotid artery abnormality	0
Systolic blood pressure difference in arms ≥20 mmHg	0
Additional imaging criteria	

Number of affected arterial territories (select one)	
• One arterial territory	0
• Two arterial territories	0
• Three or more arterial territories	+3
Symmetric involvement of paired arteries [‡]	+1
Abdominal aorta involvement with renal or mesenteric involvement [‡]	+3
SUM:	10

References

1 Grayson PC, Ponte C, Suppiah R, et al; DCVAS Study Group. 2022 American College of Rheumatology/EULAR classification criteria for Takayasu arteritis. *Ann Rheum Dis.* 2022; 81: 1654-1660.