## Supplementary material

Cylke-Falkowska K, Czajka-Oraniec I, Stelmachowska-Banaś M, Zgliczyński W. Immune checkpoint inhibitor—induced central diabetes insipidus in the course of panhypophysitis. Pol Arch Intern Med. 2024; 134: 16672. doi:10.20452/pamw.16672

Please note that the journal is not responsible for the scientific accuracy or functionality of any supplementary material submitted by the authors. Any queries (except missing content) should be directed to the corresponding author of the article.

Table S1

Hormonal test	Baseline results	Follow-up (4months)	Follow-up (15 months)	Normal range
Cortisol [ug/dl]	1.0	2.0	< 0.5	6-18.4
ACTH [pg/ml]	3.1	1.6	2.3	6-46
DHEA-S [ug/dl]	7	6	5	97-392
TSH [mIU/l	0.004	0.005	0.564	0.35-4
fT4 [pmol/l]	8.96	15.15	10.62*	10.29-22.7
GH [ug/l]	< 0.05	< 0.05	< 0.05	4.04-15.2
LH [mIU/ml]	0.3	0.4	< 0.2	2.15-2.6
FSH [IU/l]	1.5	0.9	0.4	1.3-11.8
PRL [ng/ml]	< 0.5	2.0	< 0.5	4.04-15.2
TEST [ng/ml]	< 0.07	0.07	5.48**	1.75-7.81
IGF-1 [ng/ml]	167	91.1	72	72-225

<sup>\*-</sup> during L-thyroxin replacement therapy

<sup>\*\*-</sup> during testosterone replacement therapy