Supplementary material

Suligowska K, Czarniak P, Pakalska-Korcala A, et al. The necessity of prevention and treatment in the population of adolescents based on a comprehensive health assessment: the SOPKARD-Junior survey. Pol Arch Intern Med. 2020; 130: 358-367. doi:10.20452/pamw.15280

Please note that the journal is not responsible for the scientific accuracy or functionality of any supplementary material submitted by the authors. Any queries (except missing content) should be directed to the corresponding author of the article.

Table S1. Percentage (%) of children at high risk of developing a disease entity (Group 2)

 according to ICD classification.

	ICD		Percentage of
	classificatio		children [% (n)]
Group 2	n	Disease entity	
electrocardiographic	Q 21.1	Atrial Septal Defect	0.75 (1)
electrocar	Q 25.8	Other congenital malformations of great arteries	0.75 (1)
graphic and	Q 23.1	Congenital insufficiency of aortic valve (Bicuspid aortic valve)	0.75 (1)
Echocardiographic assessment	Q 22.8	Other congenital malformations of tricuspid valve	0.75 (1)
Orthopaedic assessment	M21.4	Flat foot [pes planus] (acquired)	30.3(30)

		ICD		Percentage of
Grou	р 3	classification	Disease entity	children [% (n)]
Echocardiographic and electrocardiographic assessment	Q 21.1	Atrial Septal Defect	0.75 (1)	
	Q 23.1	Congenital insufficiency of aortic valve	3.0 (4)	
	Q 23.3	Congenital mitral insufficiency	6.0 (8)	
	I49.1	Atrial premature depolarization (Atrial premature beats)	0.75 (1)	
		M21.4	Flat foot [pes planus](acquired)	12.1 (12)
		M92.5	Juvenile osteochondrosis of tibia and fibula	2.0 (2)
		M40.1	Other secondary kyphosis	2.0 (2)
		Q66.2	Metatarsus varus	6.0(6)
Orthopaedic assessment	ent	Q68.4	Congenital bowing of tibia and fibula	1.0(1)
	dic assessm	Q65.8	Other congenital deformities of hip (Femoral torsion)	3.0 (3)
	Orthopae	M20.1	Hallux valgus (acquired)	2.0 (2)

Table S2. Percentage (%) of children with a detected or reported disease entity according toICD classification.