

Supplementary material

Fleischer-Stępniewska K, Rymer W, Inglot MS, et al. Risk of autoimmune hepatitis reactivation in patients with chronic hepatitis C and autoimmune hepatitis treated with direct-acting antivirals. Pol Arch Intern Med. 2019; 129: 215-218. doi:10.20452/pamw.4442

Please note that the journal is not responsible for the scientific accuracy or functionality of any supplementary material submitted by the authors. Any queries (except missing content) should be directed to the corresponding author of the article.

Table S1. Immune parameters during antiviral therapy and at follow-up

Patient no.	Visit	ANA antinuclear		ANA anticytoplasmic [titer]	ASMA [titer]	immunoglobulin G [xUNL]	□-globulin [xUNL]	AIH therapy
		pattern [§]	titer					
1	D0	-h	1:320	1:100	undetected	1.64	1.98	none
	EOT	-h	1:320	undetected	0.11	1.8	1.6	none
	FU24	ND		ND	ND	1.54	1.6	PDN/AZA ¹
	FU2Y	-h	1:320	1:100	undetected	1,29	1.48	PDN/AZA
2	D0	-s	1:100	undetected	undetected	0.77	0.86	BUD/AZA ²
	EOT	-fd	1:320	undetected	ND	0.5	0.67	PDN/AZA
	FU24	-fd	1:320	undetected	undetected	0.55	ND	PDN/AZA
	FU2Y	-fd	1:100	undetected	undetected	0,52	0,65	PDN/AZA

3	D0	-n	1:320	1:100	undetected	ND	0.85	BUD/AZA ³
	EOT	-n	1:100	undetected	undetected	0.57	0.55	BUD/AZA
	FU24	-n	1:100	1:100	undetected	0.59	0.68	BUD/AZA
	FU2Y	-n	1:320	1:100	undetected	0,60	0,70	none
4	D0	-		1:3200	1:320	1.1	1.36	none
	EOT	-s	1:100	1:1000	1:1000	1.0	1.06	none
	FU24	-s	1:100	undetected	undetected	1.1	1.10	none
	FU2Y	-s	1:100	1:320	1:320	1,0	1,02	none

^sANA-antinuclear pattern: -h – homogeneous, -fd – few dots, -s – speckled, -n – nucleolar;

ND – not done; UNL – upper norm level; PDN – prednisone; AZA – azathioprine; BUD – budesonide;

1 – at 9th week of therapy started with 40mg PDN, then maintenance therapy: 10mg of PDN with 100 mg QD (once a day) of AZA; continued

2 – before DAA therapy: 3mg TID (three times a day) of BUD with 50mg QD of AZA; at baseline: 3mg QD of BUD with 50mg QD of AZA, then modified schedule of corticosteroid to 3mg BUD every second day, and finally to 5mg QD of PDN instead of BUD; continued

3 – before DAA therapy: 3mg QD of BUD with 25 of AZA; at baseline 3mg of BUD every second day with 25 of AZA with; modified schedule from FU24 to withdraw of immunosuppressive therapy at 48 week of follow-up DAA therapy

