

Supplementary material

Janik MK, Wunsch Ewa, Moskwa M, et al. Depression in patients with autoimmune hepatitis: the need for detailed psychiatric assessment. *Pol Arch Intern Med.* 2019; 129: 645-647. doi:10.20452/pamw.14898

Please note that the journal is not responsible for the scientific accuracy or functionality of any supplementary material submitted by the authors. Any queries (except missing content) should be directed to the corresponding author of the article.

Supplementary material

Table S1. Clinical characteristics and results of the tests in included patients at psychiatric assessment .

CHARACTERISTIC	RESULT
MALES / FEMALES	2 (20%) / 8 (80%)
AGE (years) mean (SD) [range]	39.1 (15.6) [19-69]
BMI (kg/m²) mean (SD)	26.4 (5.9)
ALT (IU/ml) mean (SD)	25 (12)
LSM (kPa) mean (SD)	9 (4)
CIRRHOSIS Y/N	2 (20%) / 8 (80%)
GAD-7 mean (SD) [range]	7.0 (6.1) [1-21]
SIPAT mean (SD) [range]	15.6 (8.2) [4-31]
AUDIT mean (SD) [range]	1.4 (1.9) [0-5]
TREATMENT	
PRED + AZA	4 (40%)
BUDE + AZA	3 (30%)
AZA	3 (30%)

The data was presented as mean values, standard deviation (SD) and ranges, when appropriated. Abbreviations: ALT, alanine aminotransferase; AUDIT, Alcohol Use Disorders Identification Test; AZA, Azathioprine; BMI, Body Mass Index; BUDE, Budesonide; GAD-7, Generalized Anxiety Disorder 7; LSM, liver stiffness measurement; PRED, Prednisolone; SIPAT, Stanford Integrated Psychosocial Assessment for Transplant.